

Lactamator



THE PRODUCT

LACTAMATOR™ is an innovative enzyme-based product developed by CPC Biotech specifically designed for the inactivation of a wide range of beta-lactam antibiotics. **LACTAMATOR™** efficiently inactivates penicillins, cephalosporins of first, second, third, fourth, fifth generation, penems and monobactams.

FORMULATION AND PACK SIZE

Freeze dried **LACTAMATOR™** is available in the following pack size:

- cod. LCT-5: pack size of 10 vials each of which contains > 50 IU of cephalosporinase activity/vial and > 500 IU of penicillinase activity/vial
 - cod. LCT-20: pack size of 1 vial containing > 2,000 IU of cephalosporinase activity/vial and > 20,000 IU of penicillinase activity/vial
 - cod. LCT-BULK (on demand)

Sterile freeze dried formulation:

- cod. LCT-5 IR: pack size of 10 vials each of which contains > 50 IU of cephalosporinase activity/vial and > 500 IU of penicillinase activity/vial
- The product is sterile by β -irradiation.

Freeze dried and sterile freeze dried **LACTAMATOR™** are also available as enzymatic products having only cephalosporinase activity (product on demand).

Sterile liquid ready to use LACTAMATOR™ is available in the following pack size:

- cod. LCT-LQS10: pack of 5 x 10 ml vials each of which contains > 100 IU of cephalosporinase activity/vial and > 1,000 IU of penicillinase activity/vial

1 IU is defined as the amount of enzyme needed to hydrolyze 1 μ mole of Penicillin G (Penicillinase) or 1 μ mole of Cephalosporin C (Cephalosporinase) per minute at 25°C and pH 7.0.

1 IU of Penicillinase corresponds to 600 Levy Units.

APPLICATION

LACTAMATOR™ is a product specifically designed for the inactivation of a broad range of beta-lactam antibiotics, thus find its application in beta-lactams sterility testing (either environmental and antibiotics sterility testing).

LACTAMATOR™ can be used for the inactivation of beta-lactams from blood or tissue sample prior to routine microbiological examination. Due to its very broad specificity range **LACTAMATOR™** can also be used in the assessment of the susceptibility of new beta-lactams antibiotics to inactivation by ESB lactamases.



USAGE

Freeze dried formulation

cod. LCT-5/LCT-5 IR: reconstitute each vial with 5 ml of purified water obtaining a solution having > 10 IU/ml of cephalosporinase activity and > 100 IU of penicillinase activity.

cod. LCT-20: reconstitute the vial with 20 ml of purified water obtaining a solution having > 100 IU/ml of cephalosporinase activity and > 1.000 IU/ml of penicillinase activity.

After reconstitution sterile filter the solution using a suitable device. Add an appropriate amount of enzyme solution to the test sample. The amount of **LACTAMATOR™** to be added to the test sample should be determined and set-up case by case depending on the application, concentration of antibiotic that should be inactivated, and depending on the specific beta-lactam that should be inactivated.

Sterile liquid ready to use formulation (cod. LCT-LQS10)

Sterile liquid **LACTAMATOR™** is an optimized ready to use solution that can be directly added to the test samples.

The amount of liquid sterile **LACTAMATOR™** to be added to the test sample should be determined and set-up case by case depending on the application, concentration of antibiotic that should be inactivated, and depending on the specific beta-lactam that should be inactivated.

STABILITY OF REAGENTS

Freeze dried formulation

Freeze dried **LACTAMATOR™** has a stability of 36 months from manufacturing date. After reconstitution and sterile filtration, the solution can be stored at -25°C/+8°C for at least three months without losing activity.

Sterile liquid ready to use formulation (cod. LCT-LQS10)

Sterile liquid **LACTAMATOR™** has a stability of 48 months from manufacturing date. The product should be stored at +2°C/+8°C

ADDITIONAL INFORMATION

LACTAMATOR™ DOES NOT CONTAIN ANIMAL DERIVATIVES and it's certified as a TSE/BSE free product.



CPC Biotech S.r.l.

Office: Via L. Galvani, 1 | 20875 | Burago di Molgora (Monza Brianza) Italy

T: +39.039.6898029 | F: +39.039.6899774 | info@cpcbiotech.it | www.cpcbiotech.it